



RM 108 | GDU 2650 3220A E-Z Vip AVAILABLE RIB: YES


Management & Positioning

- Best suited for drought stressed to moderate yield environments
- Widely adapted, best west and central
- Good plant health in general and good tolerance for Goss's wilt
- Adapted to a wide population range with semi-flex ear type
- Medium plant stature, very good early vigor

Precision Placement™ Management

Planting Date		Soils	
Early	HR	Clay Loams	R
Late	R	Sandy	R
Variable Planting Populations		Silt Loam	HR
With Yield Zone		Peat	R
Low	24-28,000	Compacted	N
Moderate	26-30,000	Poorly Drained	N
High	32-38,000	Drought Prone	R
Very High	N	High pH	R
Dryland <20	16-24,000	Fertility	
<i>Population=(Yield Goal/7.5)*1000</i>		Nitrogen	
Water Management		Low	N
Full Irrigation	R	Med	R
Limited	R	High	R
Dryland	HR	Post Application	
Crop Rotation		Herbicide	Normal
Corn/Soybeans	HR	Fungicide	Positive
Continue Corn	w/Fungicide	LPI Nutritional	Very Good
Tillage		Herbicide Resistance	Glyph / Gluf
Conventional	HR	Harvest Schedule	
Minimum	HR	Early	HR
Ridge-Till	HR	Late	R
No-Till	R	Forage / Silage Quality	
Soil Productivity		Silage Select	N
Low	R	Dual Purpose	R
Moderate	HR		
High	R		

Agronomic Ratings



Agronomic Traits

Plant Height	Medium	Kernel Rows	14-18
Ear Height	Medium-Low	Cob Color	Pink
Flowering	Medium	Kernel Texture	Medium
Leaf Habit	Semi-Upright	Kernel Depth	Medium
Ear Flex	Semi-Flex	Husk Coverage	Adequate
Ear Type	Medium	Shank Length	Medium

Disease Tolerance Ratings

Gray Leaf Spot	7	Common Rust	6
Goss's Wilt	8	Southern Rust	5
N. Leaf Blight	6	Anthracnose	6
S. Leaf Blight	6	L. Anthracnose	n/a
Eye Spot	n/a		

Trait Versions Available

CONV - NONE

Plant with These Hybrids for Diversity

D49VC70 | D50VC78 | D50VC09 | D51VC41

Ratings Key: 9=Excellent, 5=Average, 1=Poor; HR=Highly Recommended, R=Recommended, N=Not Recommended, n/a Testing not complete. Herbicide abbreviations: GR=Growth Regulator, PI=Pigment Inhibitor, SU=Sulfonylurea. Yield zones based upon yield goals in field.
 Actual ratings based on best current information available and may be affected by changing environmental and management conditions.